arXiv:1205.1592v2 [math.GT] 20 Oct 2012

# Singular Fibers in Barking Families of Degenerations of Elliptic Curves

Takayuki OKUDA

**Abstract**

Takamura [Ta3] established a theory on splitting families of degenerations of complex curves. He introduced a powerful method for constructing a splitting family, called a *barking family*, in which there appear not only a singular fiber over the origin but also singular fibers over other points, called subordinate fibers. In this paper, for the case of degenerations of elliptic curves, we determine the types of these subordinate fibers.

## 1 Introduction

Let $\pi : M \to \Delta$ be a proper surjective holomorphic map from a smooth complex surface $M$ to an open disk $\Delta := \{s \in \mathbb{C} \ : \ |s| < \delta\}$ $(\delta > 0)$ on $\mathbb{C}$. We call $\pi : M \to \Delta$ a *family of complex curves* of genus $g \geq 1$ if all but a finite number of fibers of $\pi$ are smooth complex curves of genus $g$. In particular, $\pi : M \to \Delta$ is called a *degeneration* (or *degenerating family*) of complex curves of genus $g$ if the fiber $X_0 := \pi^{-1}(0)$ over the origin is singular and the other fibers $X_s := \pi^{-1}(s)$ $(s \neq 0)$ are smooth.

In this paper, we are interested in the problem: *How does the singular fiber split in a deformation?* Let us recall the concept of splitting families of degenerations. Let $\mathcal{M}$ be a smooth complex 3-manifold and set $\Delta^\dagger := \{t \in \mathbb{C} \ : \ |t| < \epsilon\}$ $(\epsilon > 0)$, a sufficiently small disk. Consider a proper flat surjective holomorphic map $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$. For any $t \in \Delta^\dagger$, set $\Delta_t := \Delta \times \{t\}$, $M_t := \Psi^{-1}(\Delta_t)$ and $\pi_t := \Psi\big|_{M_t} : M_t \to \Delta_t$. Suppose that $\pi_0 : M_0 \to \Delta_0$ coincides with a given degeneration $\pi : M \to \Delta$. Then we call $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$ a *deformation family* of the degeneration $\pi : M \to \Delta$. In particular, $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$ is called a *splitting family* if for any $t \in \Delta^\dagger \setminus \{0\}$, $\pi_t : M_t \to \Delta_t$ is a family of complex curves with at least two singular fibers. In this case, setting $X_{s,t} := \Psi^{-1}(s,t)$, we say that the original singular fiber $X_0$ in $M$ (that is, $X_{0,0}$ in $M_0$) *splits into* singular fibers $X_{s_1,t}, X_{s_2,t}, \ldots, X_{s_N,t}$ in $M_t$ $(t \neq 0)$.

---

2010 *Mathematics Subject Classification.* Primary 14D06, 32S50; Secondary 14H15, 14D05.

*Key words and phrases.* Degeneration of complex curves, Splliting family, Elliptic curve, Singular fiber, Monodromy.

To classify *atomic degenerations* — degenerations admitting no splitting family — Takamura [Ta3] gave a powerful method for constructing splitting families. Splitting families obtained by this construction are called *barking families.* In a barking family, the original singular fiber $X_0$ splits into a singular fiber over the origin (the *main fiber*) and other singular fibers (*subordinate fibers*). Takamura describes the main fiber explicitly in such a way that a part of the original singular fiber $X_0$ looks "barked" (peeled) off from $X_0$. See Figure 2 in Section 2.

In this paper, for the case of degenerations of elliptic curves (complex curves of genus 1), we will almost completely describe the subordinate fibers appearing in barking families. In Section 2, we first review Takamura's theory on barking deformations, mainly for degenerations with *stellar* (star-shaped) singular fibers. In fact, most of the degenerations of elliptic curves may have stellar singular fibers. In Section 3, we will give the main theorem — a list of barking deformations, which describes the types of singular fibers appearing in them. To obtain the list, in Section 4, we determine the types of subordinate fibers. We focus on the sum of the Euler characteristics of singular fibers and give a list of combinations of subordinate fibers that can appear for each deformation. Then we determine their types (except for some deformations[1]) in two ways: (1) by compairing the traces of their monodromies and (2) by investigating the number of the subordinate fibers and that of their singularities.

**Acknowledgements** The author would like to express his deep gratitude to Shigeru Takamura for fruitful discussions. The author would also like to thank Masaaki Ue and Osamu Saeki for helpful comments and suggestions.

## 2 Takamura's theory

In this section, we review Takamura's theory on barking deformations. For details see [Ta3].

First we shall recall the concept of linear degenerations. We begin with preparation. Let $\pi : M \to \Delta$ be a degeneration of complex curves of genus $g \geq 1$ and express its singular fiber as $X_0 = \sum_i m_i \Theta_i$, where $\Theta_i$ is an irreducible component of $X_0$ with multiplicity $m_i$. We call $\pi : M \to \Delta$ (or $X_0$) is *normally minimal* if the following conditions are satisfied: (i) Any singularity of the *underlying reduced curve* $X_{0,red} := \sum_i \Theta_i$ of $X_0$ is a node. (ii) If $\Theta_i$ is a $(-1)$-curve (an exceptional curve of the first kind), then $\Theta_i$ intersects other irreducible components at at least three points. In what follows, we assume that any given degeneration $\pi : M \to \Delta$ is normally minimal[2],

---

[1] See (4.8), (5.3), (5,4) and (7.2) in the list of Theorem 3.1.

[2] In [Ta2] and [Ta3], we define linear degenerations for the case that $X_{0,red}$ has at most *simple normal crossings*, that is, (i) any singularity of $X_{0,red}$ is a node and (iii) any irreducible component $\Theta_i$ is not self-intersecting. Note that all normally minimal degenerations of elliptic curves but of type $mI_1$ satisfy the condition (iii).

For an irreducible component $\Theta_i$ of $X_0$, we denote by $N_i$ the normal bundle of $\Theta_i$ in $M$. Let $\{p_i^{(1)}, p_i^{(2)}, \dots p_i^{(h)}\}$ be the set of the intersection points on $\Theta_i$ with other irreducible components of $X_0$ and $m^{(j)}\,(j = 1, 2, \dots, h)$ be the mulitiplicity of the irreducible component intersecting $\Theta_i$ at $p_i^{(j)}$. Then there exists a holomorphic section $\sigma_i$ of the line bundle $N_i^{\otimes(-m_i)}$ such that

$$\mathrm{div}(\sigma_i) = \sum_{j=1}^{h} m^{(j)} p_i^{(j)},$$

where $\mathrm{div}(\sigma_i)$ denotes the divisor defined by $\sigma_i$. Here $\sigma_i$ has a zero of order $m^{(j)}$ at $p_i^{(j)}$. Note that $\sigma_i$ is uniquely determined up to multiplication by a constant. We call $\sigma_i$ the *standard section* of $N_i^{\otimes(-m_i)}$ on $\Theta_i$.

Take an open covering $\Theta_i = \bigcup_\alpha U_\alpha$ such that $U_\alpha \times \mathbb{C}$ is a local trivialization of the normal bundle $N_i$ on $\Theta_i$. We denote by $(z_\alpha, \zeta_\alpha)$ coordinates of $U_\alpha \times \mathbb{C}$. Now define a holomorphic function $\pi_{i,\alpha} : U_\alpha \times \mathbb{C} \to \mathbb{C}$ by

$$\pi_{i,\alpha}(z_\alpha, \zeta_\alpha) := \sigma_{i,\alpha}(z_\alpha)\zeta_\alpha^{m_i},$$

where $\sigma_{i,\alpha}$ is the local expression of $\sigma_i$ on $U_\alpha$. Then the set $\{\pi_{i,\alpha}\}_\alpha$ of holomorphic functions defines a global holomorphic function $\pi_i : N_i \to \mathbb{C}$.

**Definition 2.1.** A degeneration $\pi : M \to \Delta$ is called *linear* if for any irreducible component $\Theta_i$ of its singular fiber $X_0$,

(1) a tubular neighborhood $N(\Theta_i)$ of $\Theta_i$ in $M$ is biholomorphic to a tubular neighborhood of a zero-section of the normal bundle $N_i$, and

(2) under the identification by the biholomorphism of (1), the following conditions are satisfied:

   (a) The restriction $\pi\big|_{N(\Theta_i)}$ coincides with the holomorphic function $\pi_i$ defined above.

   (b) If $\Theta_i$ intersects $\Theta_j$ at a point $p$, then there exist local trivializations $U_\alpha \times \mathbb{C}$ of $N_i$ and $U_\beta \times \mathbb{C}$ of $N_j$ around $p$ such that neighborhoods of $p$ in $N(\Theta_i)$ and $N(\Theta_j)$ are identified by *plumbing* $(z_\alpha, \zeta_\alpha) = (\zeta_\beta, z_\beta)$ and $\pi$ is locally expressed as

$$\pi\big|_{N(\Theta_i)}(z_\alpha, \zeta_\alpha) = z_\alpha^{m_j}\zeta_\alpha^{m_i} \quad \textit{and} \quad \pi\big|_{N(\Theta_j)}(z_\beta, \zeta_\beta) = z_\beta^{m_i}\zeta_\beta^{m_j},$$

   where $(z_\alpha, \zeta_\alpha) \in U_\alpha \times \mathbb{C}$ and $(z_\beta, \zeta_\beta) \in U_\beta \times \mathbb{C}$.

**Remark 2.2.** Any degeneration of complex curves (possibly, not normally minimal), after successive blowing up and down, becomes a degeneration toplogically equivalent to some linear degeneration.



Figure 1: A sigular fiber of type $II^*$ of a degeneration of elliptic curves is stellar. Each circle denotes a projective line, the number stands for its multiplicity, and each intersection point is a node.

If $\pi : M \to \Delta$ is linear, then we may express $M$ locally as a hypersurface in some space as follows: We first identify $M$ with the graph of $\pi$ in $M \times \Delta$

$$\mathrm{Graph}(\pi) = \{(x, s) \in M \times \Delta \,:\, \pi(x) - s = 0\}$$

via the natural projection $\mathrm{Graph}(\pi) \ni (x, s) \to x \in M$. Recall that for any irreducible component $\Theta_i$ of the singular fiber $X_0$, the map $\pi$ is expressed around $\Theta_i$ as

$$\pi(z_i, \zeta_i) = \sigma_i(z_i)\zeta_i^{m_i},$$

where $\sigma_i$ is the standard section of $N_i^{\otimes(-m_i)}$ on $\Theta_i$. Then we obtain the local expression of $M$ around $\Theta_i$:

$$H_i : \quad \sigma_i(z_i)\zeta_i^{m_i} - s = 0 \quad \text{in } N_i \times \Delta. \tag{2.1}$$

Note that these hypersurfaces are glued around their intersection points by plumbing $(z_j, \zeta_j, s) = (\zeta_i, z_i, s)$ where $(z_i, \zeta_i, s) \in N_i \times \Delta$ and $(z_j, \zeta_j, s) \in N_j \times \Delta$.

For a linear degeneration $\pi : M \to \Delta$, its singular fiber $X_0$ consists of three kinds of parts: *cores*, *branches* and *trunks*. A irreducible component $\Theta_i$ of $X_0$ is called a *core* if (a) $\Theta_i$ intersects other irreducible components at at least three points or (b) $\Theta_i$ is not a projective line. A *branch* is a chain $\sum_i m_i\Theta_i$ of projective lines attached with a core on one hand, while a *trunk* is a chain $\sum_i m_i\Theta_i$ of projective lines connecting other irreducible components on both hands. We say that $X_0$ is a *stellar* singular fiber if $X_0$ consists of a core and branches emanating from it. See Figure 1. Otherwise $X_0$ is said to be *constellar*.

We obtain a 'degeneration' whose singular fiber is a *fringed* branch of length $\lambda$ as follow: Let $m_0, m_1, \dots, m_{\lambda+1}$[3] ($\lambda \geq 1$) be nonnegative integers such that

$$\begin{cases} m_0 > m_1 > \cdots > m_\lambda > m_{\lambda+1} = 0 \text{ and} \\ r_i := (m_{i-1} + m_{i+1})/m_i \text{ is an integer greater than 1 } (i = 1, 2, \dots, \lambda). \end{cases} \tag{\#}$$

---

[3] In this paper, by convention, we append $m_{\lambda+1} = 0$ to the positive integers $m_0, m_1, \dots, m_\lambda$ (see [Ta3] Section 5.1, p.85), so that $r_\lambda := (m_{\lambda-1} + m_{\lambda+1})/m_\lambda$ equals $m_{\lambda-1}/m_\lambda$.

We take $\lambda$ copies $\Theta_1, \Theta_2, \ldots, \Theta_\lambda$ of the projective line. Let $\Theta_i = U_i \cup V_i$ be a open covering by two complex lines $U_i, V_i$ $(= \mathbb{C})$ with coordinates $w_i \in U_i$ and $z_i \in V_i$ satisfying $z_i = 1/w_i$. Next we obtain a line bundle $N_i$ on $\Theta_i$ of order $-r_i$ from $U_i \times \mathbb{C}$ and $V_i \times \mathbb{C}$ by identifying $(z_i, \zeta_i) \in V_i \times \mathbb{C}$ with $(w_i, \eta_i) \in U_i \times \mathbb{C}$ via

$$g_i \, : \, z_i = \frac{1}{w_i}, \quad \zeta_i = w_i^{r_i} \eta_i.$$

Now consider the hypersurface $W_i$ in $N_i \times \Delta$ defined by

$$\begin{cases} H_i : w_i^{m_{i-1}} \eta_i^{m_i} - s = 0, & \text{in } U_i \times \mathbb{C} \times \Delta, \\ H_i' : z_i^{m_{i+1}} \zeta_i^{m_i} - s = 0, & \text{in } V_i \times \mathbb{C} \times \Delta. \end{cases} \tag{2.2}$$

Under the plumbings $(w_{i+1}, \eta_{i+1}, s) = (\zeta_i, z_i, s)$ of $N_i$ and $N_{i+1}$ $(i = 1, 2, \ldots, \lambda - 1)$, the hypersurfaces $W_1, W_2, \ldots, W_\lambda$ are glued, so that they together define a smooth complex surface $M$. Letting $\pi : M \to \Delta$ be the natural projection, the central fiber is

$$\pi^{-1}(0) = m_0 \mathbb{C} + m_1 \Theta_1 + m_2 \Theta_2 + \cdots + m_\lambda \Theta_\lambda.$$

Thus:

**Lemma 2.3.** *Let $m_0, m_1, \ldots, m_{\lambda+1}$ $(\lambda \geq 1)$ be integers satisfying the following conditions:*

$$\begin{cases} m_0 > m_1 > \cdots > m_\lambda > m_{\lambda+1} = 0. \\ r_i := (m_{i-1} + m_{i+1})/m_i \text{ is an integer greater than } 1 \ (i = 1, 2, \ldots, \lambda). \end{cases} \tag{\#}$$

*Then there exists a degeneration with the singular fiber*

$$X_0 = m_0 \Delta_0 + m_1 \Theta_1 + m_2 \Theta_2 + \cdots + m_\lambda \Theta_\lambda,$$

*where $\Delta_0$ is an open disk and $\Theta_1, \Theta_2, \ldots, \Theta_\lambda$ are projective lines such that each pair of $\Theta_i$ and $\Theta_{i+1}$ $(i = 1, 2, \ldots, \lambda - 1)$ and $\Delta_0$ and $\Theta_1$ intersect transversely at one point.*

Next we define a special subdivisor of stellar singular fibers. Let $\pi : M \to \Delta$ be a linear degeneration of complex curves with the stellar singular fiber $X_0 = m_0 \Theta_0 + \sum_{j=1}^h \mathrm{Br}^{(j)}$, where $\Theta_0$ is the core and $\mathrm{Br}^{(j)}$ $(j = 1, 2, \ldots, h)$ is a branch. Express as $\mathrm{Br}^{(j)} = m_1^{(j)} \Theta_1^{(j)} + m_2^{(j)} \Theta_2^{(j)} + \cdots + m_{\lambda^{(j)}}^{(j)} \Theta_{\lambda^{(j)}}^{(j)}$ and let $\overline{\mathrm{Br}}^{(j)} = m_0 \Delta_0^{(j)} + m_1^{(j)} \Theta_1^{(j)} + \cdots + m_{\lambda^{(j)}}^{(j)} \Theta_{\lambda^{(j)}}^{(j)}$ be a fringed branch. Consider a connected subdivisor $Y = n_0 \Theta_0 + \sum_{j=1}^h \mathrm{br}^{(j)}$ of $X_0$, where $\mathrm{br}^{(j)} := n_1^{(j)} \Theta_1^{(j)} + n_2^{(j)} \Theta_2^{(j)} + \cdots + n_{\nu^{(j)}}^{(j)} \Theta_{\nu^{(j)}}^{(j)}$ $(j = 1, 2, \ldots, h)$. Here $Y$ satisfies $0 \leq \nu^{(j)} \leq \lambda^{(j)}$ and $0 < n_i^{(j)} \leq m_i^{(j)}$ for each $i$ and $j$. Set $\overline{\mathrm{br}}^{(j)} := n_0 \Delta_0^{(j)} + n_1^{(j)} \Theta_1^{(j)} + n_2^{(j)} \Theta_2^{(j)} + \cdots + n_{\nu^{(j)}}^{(j)} \Theta_{\nu^{(j)}}^{(j)}$. For the time being, we focus on $\overline{\mathrm{Br}}^{(j)}$ and $\overline{\mathrm{br}}^{(j)}$, omitting the superscript $(j)$ to simplify notation. We call $\overline{\mathrm{br}}$ a *subbranch* of $\overline{\mathrm{Br}}$ if one of the following conditions is satisfied:

(1) $\nu = 0, 1$, or (2) $\nu \geq 2$ and $n_{i+1} = r_i n_i - n_{i-1}$ $(i = 1, 2, \ldots, \nu - 1)$,

where $r_i := (m_{i-1} + m_{i+1})/m_i$ (see (#) in Lemma 2.3). Inductively set $n_{\nu+1} := r_\nu n_\nu - n_{\nu-1}$ $(\nu \geq 1)$. If $\nu = 0$, then we set $n_{\nu+1} = n_1 := 0$. Next define the three types of subbranches for a positive integer $l$ as follows:

**Type $A_l$.** A subbranch $\overline{\mathrm{br}}$ of $\overline{\mathrm{Br}}$ is of *type* $A_l$ if $ln_i \leq m_i$ for each $i$ and $n_{\nu+1} \leq 0$.

**Type $B_l$.** A subbranch $\overline{\mathrm{br}}$ of $\overline{\mathrm{Br}}$ is of *type* $B_l$ if $ln_i \leq m_i$ for each $i$, $n_\nu = 1$ and $m_\nu = l$.

**Type $C_l$.** A subbranch $\overline{\mathrm{br}}$ of $\overline{\mathrm{Br}}$ is of *type* $C_l$ if $ln_i \leq m_i$ for each $i$, $n_\nu = n_{\nu+1}$ and $m_\nu - m_{\nu+1}$ divides $l$.

Now we return to a connected subdivisor $Y$ of the stellar singular fiber $X_0$.

**Definition 2.4.** Let $Y = n_0\Theta_0 + \sum_{j=1}^{h} \mathrm{br}^{(j)}$ be a connected subdivisor of $X_0$ such that $n_0 < m_0$ and each $\overline{\mathrm{br}}^{(j)}$ is a subbranch of $\overline{\mathrm{Br}}^{(j)}$. Then $Y$ is called a *crust* of $X_0$ if there exists a meromorphic section $\tau$ of the line bundle $N_0^{\otimes n_0}$ on $\Theta_0$ such that for some nonnegative divisor $D = \sum_{i=1}^{k} a_i q_i$ on $\Theta_0$,

$$\mathrm{div}(\tau) = -\sum_{j=1}^{h} n_1^{(j)} p^{(j)} + D,$$

where $N_0$ denotes the normal bundle of $\Theta_0$ in $M$, $\{p^{(j)}\}$ is the set of the attachment points on $\Theta_0$ with the branches $\mathrm{Br}^{(j)}$. Moreover, for a positive integer $l$, if each $\overline{\mathrm{br}}^{(j)}$ is a subbranch of $\overline{\mathrm{Br}}^{(j)}$ of either type $A_l$, type $B_l$ or type $C_l$, then we call $Y$ a *simple crust* of $X_0$ with *barking multiplicity* $l$.

We call the meromorphic section $\tau$ a *core section*. Note that $\tau$ is not uniquely determined by $Y$. Setting $r_0 := \sum_{j=1}^{h} m_1^{(j)}/m_0$ and $r_0' := \sum_{j=1}^{h} n_1^{(j)}/n_0$, the following holds:

**Lemma 2.5.** *Suppose that $\Theta_0$ is a projective line. Then a connected subdivisor $Y$ is a crust of $X_0$ (equivalently, $Y$ has a core section $\tau$) if and only if $r_0 \leq r_0'$. Moreover $\tau$ has no zero, that is, $D = 0$ exactly when $r_0 \leq r_0'$.*

Now we state Takamura's theorem:

**Theorem 2.6** (Takamura [Ta3])**.** *Let $\pi : M \to \Delta$ be a linear degeneration with the stellar singular fiber $X_0$. If $X_0$ has a simple crust, then $\pi : M \to \Delta$ admits a splitting family $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$.*

Takamura constructed a deformation familly of $\pi : M \to \Delta$ associated with a simple crust $Y$, which turns out to be a splitting family. We call it a *barking family associated with* the simple crust $Y$ and this construction is called a *barking deformation.* For a singular fiber $X_{s,t} := \Psi^{-1}(s, t)$ $(t \neq 0)$ in a barking family, we say that $X_{s,t}$ is the *main fiber* if $s = 0$, and a *subordinate fiber* if $s \neq 0$. In other words, the original singular fiber splits into the main fiber and subordinate fibers. Takamura describes the main fiber $X_{0,t}$ explicitly in such a way that the simple crust $Y$ looks “barked” (peeled) off from the original singular fiber $X_0$. See Figure 2.



Figure 2: In a barking family, the singular fiber of type $II^*$ splits into the main fiber of type $III^*$ and some subordinate fiber. It seems that the simple crust is "barked" (peeled) off from the original singular fiber.

**Kodaira's notation** Before proceeding, we include Kodaira's list of singular fibers of (relatively) minimal degenerations of elliptic curves [Ko]. See Table 1 below.

For a singular fiber $X$, we denote by $e(X)$ the topological Euler characteristic of the underlying reduced curve of $X$. Let $A_X \in SL(2, \mathbb{Z})$ be the standard monodromy matrix of $X$ and $\mathrm{Tr}(A_X)$ denote the trace of $A_X$. Note that singular fibers of type $II, III, IV$ are not normally minimal. After successive blowing up, they become normally minimal. On the other hand, The other singular fibers are normally minimal.

## 3 Main result

Takamura showed the list of simple crusts (and weighted crustal sets) for barking deformations for degenerations of complex curves of genera up to 5. See Chapter 26 of [Ta3]. As we saw in Section 2, barking deformations explicitly describe the main fibers appearing in barking families. On the other hand, it is not clear what types of subordinate fibers will appear. The aim of this paper is to show the description of subordinate fibers in these barking families for the case of degenerations of elliptic curves (complex curves of genus 1).

In what follows, if the original singular fiber $X_0$ splits into the main fiber $X_{0,t}$ and subordinate fibers $X_{s_1,t}, X_{s_2,t}, \ldots, X_{s_N,t}$, then we express

$$X_0 \longrightarrow X_{0,t} \text{ with } X_{s_1,t} + X_{s_2,t} + \cdots + X_{s_N,t}$$

to distinguish $X_{0,t}$ from the others. We denote the types of singular fibers of genus 1 in Kodaira's notation. See Table 1.

Table 1: Kodaira's notation.

| | the singular fiber $X$ | $e(X)$ | $A_X$ | $\mathrm{Tr}(A_X)$ |
|---|---|---|---|---|
| $mI_0(m \geq 2)$ | a multiple torus | $0$ | $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ | $2$ |
| $mI_1$ | a (multiple) projective line<br>with one node | $1$ | $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ | $2$ |
| $mI_n(n \geq 2)$ |  | $n$ | $\begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}$ | $2$ |
| $II$ | a projective line<br>with one cusp | $2$ | $\begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}$ | $1$ |
| $III$ | two projective lines<br>with second order contact | $3$ | $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ | $0$ |
| $IV$ | three projective lines intersecting<br>transversally at one point | $4$ | $\begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$ | $-1$ |
| $I_0^*$ |  | $6$ | $\begin{pmatrix} -1 & 0 \\ -1 & 0 \end{pmatrix}$ | $-1$ |
| $I_n^*$ |  | $6+n$ | $\begin{pmatrix} -1 & -n \\ -1 & 0 \end{pmatrix}$ | $-1$ |
| $II^*$ |  | $10$ | $\begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$ | $1$ |
| $III^*$ |  | $9$ | $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ | $0$ |
| $IV^*$ |  | $8$ | $\begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix}$ | $-1$ |

**Theorem 3.1.** *For degenerations of complex curves of genus* 1 *(that is, elliptic curves), the types of subordinate fibers in the barking families listed in* [Ta3] *are presented in the following list:*

(1.1) $II \longrightarrow I_1$ with $I_1$



(1.2) $II \longrightarrow I_1$ with $I_1$



(2.1) $II^* \longrightarrow III^*$ with $I_1$



(2.2) $II^* \longrightarrow IV^*$ with $II$



### (2.3) $II^* \longrightarrow I_2^*$ with $I_1 + I_1$



### (2.4) $II^* \longrightarrow I_5$ with $I_1 + I_1 + I_1 + I_1 + I_1$



### (2.5) $II^* \longrightarrow I_3^*$ with $I_1$



### (2.6) $II^* \longrightarrow I_3^*$ with $I_1$



### (2.7) $II^* \longrightarrow I_8$ with $I_1 + I_1$



### (2.8) $II^* \longrightarrow III^*$ with $I_1$



### (2.9) $II^* \longrightarrow III^*$ with $I_1$



### (3.1) $III \longrightarrow I_2$ with $I_1$



### (3.2) $III \longrightarrow I_1$ with $I_2$



### (3.3) $III \longrightarrow I_2$ with $I_1$



### (4.1) $III^* \longrightarrow IV^*$ with $I_1$



### (4.2) $III^* \longrightarrow I_1^*$ with $I_2$



### (4.3) $III^* \longrightarrow I_2^*$ with $I_1$



(4.4) $III^* \longrightarrow I_0^*$ with $I_1 + I_1 + I_1$



(4.5) $III^* \longrightarrow I_6$ with $I_1 + I_1 + I_1$



(4.6) $III^* \longrightarrow I_2^*$ with $I_1$



(4.7) $III^* \longrightarrow I_7$ with $I_1 + I_1$



### (4.8) $III^* \longrightarrow I_6$ with $(II + I_1,\ I_2 + I_1$ or $I_1 + I_1 + I_1)$



### (4.9) $III^* \longrightarrow IV$ with $I_1$



### (5.1) $IV \longrightarrow I_3$ with $I_1$



### (5.2) $IV \longrightarrow I_2$ with $I_1 + I_1$



### (5.3) $IV \longrightarrow I_2$ with $(II$ or $I_1 + I_1)$



### (5.4) $IV \longrightarrow II$ with $(II,\ I_2$ or $I_1 + I_1)$



### (6.1) $IV^* \longrightarrow I_1^*$ with $I_1$



### (6.2) $IV^* \longrightarrow I_0^*$ with $I_1 + I_1$



(6.3) $IV^* \longrightarrow I_6$ with $I_1 + I_1$



(6.4) $IV^* \longrightarrow I_1^*$ with $I_1$



(7.1) $I_0^* \longrightarrow I_4$ with $I_1 + I_1$



(7.2) $I_0^* \longrightarrow I_3$ with $(II + I_1$ or $I_1 + I_1 + I_1)$



(8.1) $I_n^* \longrightarrow I_{n-1}^*$ with $I_1$



(8.2) $I_n^* \longrightarrow I_{n+4}$ with $I_1 + I_1$



**Remark 3.2.** (1) The list in Theorem 3.1 does not contain singular fibers of type $mI_n$. In fact, $mI_n$ splits into $mI_{n-1}$ and $I_1$ by another deformation. For details see [Ta1]. (2) In Section 2, we reviewed only barking deformations associated with one simple crust for stellar singular fibers. For details of "compounding" barking like (2.6), (2.7), (4.7) and (4.8), and barking for "constellar" singular fibers like (8.1) and (8.2), see [Ta3].

# 4 Determination of subordinate fibers

In this section, we will determine the types of subordinate fibers appearing in barking families listed in Theorem 3.1. First we recall results on Euler characteristics of singular fibers, which gives us the sum of the Euler characteristics of the subordinate fibers. And then we will approach to determination of their types in two ways: (1) by compairing the traces of their monodromies and (2) by investigating the number of the subordinate fibers and that of their singularities.

## 4.1 Euler characteristics of subordinate fibers

First we recall results on Euler characteristics of singular fibers of degenerations. We denote by $e(X)$ the (*topological*) *Euler characteristic* of the underlying reduced curve of a singular fiber $X$.

**Lemma 4.1.** *Let $\pi : M \to \Delta$ be a degeneration of complex curves of genus $g$ with the singular fiber $X_0$ and let $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$ be a splitting family of $\pi : M \to \Delta$.*
(1) *If $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$ splits $X_0$ into $X_1, X_2, \dots, X_N$ in $M_t$, then the following formula holds:*

$$e(X_0) - 2(1-g) = \sum_{i=1}^{N} \{e(X_i) - 2(1-g)\}. \tag{4.1}$$

(2) *Moreover, if $g = 1$ (for which $\pi : M \to \Delta$ is a degeneration of elliptic curves), then the following holds:*

$$e(X_0) = \sum_{i=1}^{N} e(X_i).$$

*Proof.* (1) The left hand side of (4.1) equals to the Euler characteristic $e(M)$ of $M$, while the right hand side equals to $e(M_t)$ (see p.97 of [BPV]). Since $M_t$ is deffeomorphic to $M$, we have $e(M) = e(M_t)$, which confirms the assertion. (2) follows from (1) clearly. □

Recall that for a singular fiber $X_{s,t} := \Psi^{-1}(s,t)$ ($t \neq 0$), we call $X_{s,t}$ the main fiber if $s = 0$, and a subordinate fiber if $s \neq 0$. In these terminologies we restate (2) of Lemma 4.1 as follows:

**Corollary 4.2.** *Let $\pi : M \to \Delta$ be a degeneration of elliptic curves with the singular fiber $X_0$. Suppose a splitting family $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$ of $\pi : M \to \Delta$ splits $X_0$ into the main fiber $X_{0,t}$ and subordinate fibers $X_{s_1,t}, X_{s_2,t}, \dots, X_{s_N,t}$ in $M_t$. Then the sum of the Euler characteristics of the subordinate fibers is*

$$\sum_{i=1}^{N} e(X_{s_i,t}) = e(X_0) - e(X_{0,t}).$$

Note that if $X$ is a singular fiber of a degeneration of elliptic curves except a multiple torus, then $e(X) \geq 1$ where the equality holds exactly when $X$ is a Lefschetz fiber of genus 1 (a projective line with a node). This fact together with Corollary 4.2 yields the following:

**Lemma 4.3.** *Let $\pi : M \to \Delta$ be a degenreration of elliptic curves with the singular fiber $X_0$. Suppose a splitting family $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$ of $\pi : M \to \Delta$ satisfies*

$$e(X_0) - e(X_{0,t}) = 1. \tag{4.2}$$

*Then $X_0$ splits into the main fiber $X_{0,t}$ and a subordinate fiber $X_{s,t}$ that is a Lefschetz fiber of genus 1 (a projective line with a node).*

**Remark 4.4.** Note that multiple tori (whose Euler characteristics are 0) do *not* appear as subordinate fibers. In fact, any subordinate fiber appearing in any barking family has only $A$-singularities. See Lemma 4.13.

Fortunately, many barking deformations in the list of Theorem 3.1 satisfy (4.2).

- (The case $e(X_0) - e(X_{0,t}) = 1$) The subordinate fiber appearing in each of the following deformations is $I_1$:

(1.1) $II \longrightarrow I_1$ with $I_1$

(1.2) $II \longrightarrow I_1$ with $I_1$

(2.1) $II^* \longrightarrow III^*$ with $I_1$

(2.5) $II^* \longrightarrow I_3^*$ with $I_1$

(2.6) $II^* \longrightarrow I_3^*$ with $I_1$

(2.8) $II^* \longrightarrow III^*$ with $I_1$

(2.9) $II^* \longrightarrow III^*$ with $I_1$

(3.1) $III \longrightarrow I_2$ with $I_1$

(3.3) $III \longrightarrow I_2$ with $I_1$

(4.1) $III^* \longrightarrow IV^*$ with $I_1$

(4.3) $III^* \longrightarrow I_2^*$ with $I_1$

(4.6) $III^* \longrightarrow I_2^*$ with $I_1$

(4.9) $III^* \longrightarrow IV$ with $I_1$

(5.1) $IV \longrightarrow I_3$ with $I_1$

(6.1) $IV^* \longrightarrow I_1^*$ with $I_1$

(6.4) $IV^* \longrightarrow I_1^*$ with $I_1$

(8.1) $I_n^* \longrightarrow I_{n-1}^*$ with $I_1$

If $e(X_0) - e(X_{0,t}) \geq 2$, then we need another criterion to determine the types of subordinate fibers. But by Corollary 4.2 we can narrow down candidates.

- (The case $e(X_0) - e(X_{0,t}) = 2$) The subordinate fiber appearing in each of the following deformations is one of $II$, $I_2$ and $I_1 + I_1$:

(2.2) $II^* \longrightarrow IV^*$ with ?

(2.3) $II^* \longrightarrow I_2^*$ with ?

(2.7) $II^* \longrightarrow I_8$ with ?

(3.2) $III \longrightarrow I_1$ with ?

(4.2) $III^* \longrightarrow I_1^*$ with ?

(4.7) $III^* \longrightarrow I_7$ with ?

(5.2) $IV \longrightarrow I_2$ with ?

(5.3) $IV \longrightarrow I_2$ with ?

(5.4) $IV \longrightarrow II$ with ?

(6.2) $IV^* \longrightarrow I_0^*$ with ?

(6.3) $IV^* \longrightarrow I_6$ with ?

(7.1) $I_0^* \longrightarrow I_4$ with ?

(8.2) $I_n^* \longrightarrow I_{n+4}$ with ?

- (The case $e(X_0) - e(X_{0,t}) = 3$) The subordinate fiber appearing in each of the following deformations is one of $III$, $I_3$, $II + I_1$, $I_2 + I_1$ and $I_1 + I_1 + I_1$:

(4.4) $III^* \longrightarrow I_0^*$ with ?

(4.5) $III^* \longrightarrow I_6$ with ?

(4.8) $III^* \longrightarrow I_6$ with ?

(7.2) $I_0^* \longrightarrow I_3$ with ?

- (The case $e(X_0) - e(X_{0,t}) = 5$)

$\underline{(2.4)\ II^* \longrightarrow I_5 \text{ with ?}}$

## 4.2 Monodromies around subordinate fibers

Next consider the monodromies around singular fibers appearing in splitting families.

Let $\pi : M \to \Delta$ be a (relatively) minimal degeneration of elliptic curves with the singular fiber $X_0$. We take a base point $*$ on $\Delta \setminus \{0\}$ and a loop (simple closed curve) $l$ in $\Delta \setminus \{0\}$ passing through $*$ and circuiting around the origin with a counterclockwise orientation. Then $\pi^{-1}(l)$ is a real 3-manifold and the restriction $\pi : \pi^{-1}(l) \to l$ is a $\Sigma$-bundle over $S^1$, where $\Sigma = \pi^{-1}(*)$ is a smooth fiber, that is, a elliptic curve. Here $\pi^{-1}(l)$ is obtained from $\Sigma \times [0,1]$ by the identification of the boundaries $\Sigma \times \{0\}$ and $\Sigma \times \{1\}$ via a orientation-preserving homeomorphism $h$. The isotopy class of $h$ is called the *topological monodromy* around $X_0$. Then $h$ induces an automorphism $\rho := h_*$ on $H_1(\Sigma, \mathbb{Z})$, which is called the *monodromy* around $X_0$. Under an identification of $\Sigma$ and $\mathbb{R}^2/\mathbb{Z}^2$, fixing the generators of $H_1(\Sigma, \mathbb{Z})$, we may express $\rho$ as an element of $SL(2, \mathbb{Z})$.

Next suppose that $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$ is a splitting family of the degeneration $\pi : M \to \Delta$, say, $\pi_t : M_t \to \Delta$ ($t \neq 0$) has singular fibers $X_1, X_2, \ldots, X_N$ ($N \geq 2$). Then we express $X_0 \longrightarrow X_1 + X_2 + \cdots + X_N$. Now we determine the (local) monodromy $\rho_k \in SL(2, \mathbb{Z})$ around each sigular fiber $X_k (k = 1, 2, \ldots, N)$ as follows: Set $s_k := \pi(X_k)$. We take loops $l_k$ in $\Delta \setminus \{s_1, s_2, \ldots, s_N\}$ passing through $*$ and circuiting around $s_k$ with a counterclockwise orientation. Then the loop $l_k$ induces the local monodromy $\rho_k \in SL(2, \mathbb{Z})$ around $X_k$ in $\pi_t : M_t \to \Delta$. Possibly after renumbering, we may assume that $l_1 l_2 \cdots l_N$ is homotopic to $l$

The following is known:

**Lemma 4.5.** (1) *In $SL(2, \mathbb{Z})$, each of $\rho, \rho_k$ is conjugate to the standard monodromy matrix of some type of singular fibers (see Table* 1*).*

(2)
$$\rho = \rho_1 \rho_2 \cdots \rho_N, \tag{4.3}$$
*where we take such monodromy representations that the multiplication in* (4.3) *coincides with that in $SL(2, \mathbb{Z})$.*

We prepare notation. Note that $SL(2, \mathbb{Z}) = \langle a, b \mid a^3 = b^2 = -I \rangle$ is generated by
$$a := \begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix} \quad \text{and} \quad b := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

Setting[4]
$$s_0 := a^{-1} b = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad s_2 := b a^{-1} = \begin{pmatrix} 1 & 0 \\ -1 & 0 \end{pmatrix},$$

---

[4] The notations $s_0$ and $s_2$ stand for [FM], in which $s_1 := aba$.

then $s_0$ and $s_2$ are also generators of $SL(2,\mathbb{Z})$: indeed we have $a = s_0 s_2$ and $b = s_0 s_2 s_0 = s_2 s_0 s_2$. Since $s_2 = s_0 s_2 s_0 (s_0 s_2)^{-1}$, $s_2$ is conjugate to $s_0$. Note that $s_0$ is the standard monodromy matrix of singular fibers of type $I_1$.

Next we express the standard monodromy matrices of singular fibers as a product of $s_0$ and $s_2$ as follows (see [U]):

$$\begin{array}{rcl}
I_n & : & (s_0)^n \\
II & : & s_0 s_2 \\
III & : & s_0 s_2 s_0 = s_2 s_0 s_2 \\
IV & : & s_0 s_2 s_0 s_2 \\
II^* & : & (s_0 s_2)^5 \\
III^* & : & (s_0 s_2)^4 s_0 \\
IV^* & : & (s_0 s_2)^4 \\
I_n^* & : & (s_0 s_2)^3 (s_0)^n.
\end{array}$$

The number of $s_0$ and $s_2$ of each product above coincides with the Euler characteristic of the corresponding type. Note that the singular fiber of any degeneration of elliptic curves but of type $mI_0\,(m \geq 2)$ splits into singular fibers of type $I_1$ (whose the Euler characteristic $e(I_1)$ is equal to 1) after repeating deformations, but not necessarily by once deformation. See [Ka] [M].

Now consider the decomposability of elements of $SL(2,\mathbb{Z})$. In what follows, we denote by the standard monodromy matrix of a singular fiber $X$ by $A_X$. We write $\rho^\tau = \tau\rho\tau^{-1}$, a conjugacy of $\rho$.

**Splittability of $II$**

$$\begin{array}{lcll}
II & \rightarrow & I_1 + I_1 & \qquad s_0 s_2 = A_{I_1} \cdot A_{I_1}^{s_0 s_2}
\end{array}$$

**Splittability of $III$**

$$\begin{array}{lcll}
III & \rightarrow & II + I_1 & \qquad s_0 s_2 s_0 = A_{II} \cdot A_{I_1} \\
III & \rightarrow & I_1 + I_1 + I_1 & \\
III & \rightarrow & I_2 + I_1 & \qquad s_0^2 s_0^{-1} s_2 s_0 = A_{I_2} \cdot A_{I_1}^{s_2}
\end{array}$$

**Splittability of $IV$**

$$\begin{array}{lcll}
IV & \rightarrow & II + II & \qquad (s_0 s_2 s_0)^2 = A_{II} \cdot A_{II} \\
IV & \rightarrow & II + I_1 + I_1 & \\
IV & \rightarrow & I_1 + I_1 + I_1 + I_1 & \\
IV & \rightarrow & I_2 + II & \qquad s_0^2 s_2 (s_0 s_2) s_2^{-1} = A_{I_2} \cdot A_{II}^{s_2} \\
IV & \rightarrow & I_2 + I_1 + I_1 & \\
IV & \rightarrow & III + I_1 & \qquad s_0 s_2 s_0 s_2 = A_{III} \cdot A_{I_1}^{s_0 s_2} \\
IV & \rightarrow & I_3 + I_1 & \qquad s_0^3 s_0^{-1} s_2 s_0 = A_{III} \cdot A_{I_1}^{s_2}
\end{array}$$

**Proposition 4.6.** *$IV$ does not admit the following decomposition:*

$$IV \longrightarrow I_2 + I_2$$

*Proof.* If $IV$ splits into $I_2 + X$, then for a monodromy matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ of $X$,

$$B := \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} A$$

is a monodromy matrix of $II^*$. Accordingly $B$ is conjugate to $\begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$. Denoting by $\mathrm{Tr}(A)$ (resp. $\mathrm{Tr}(B)$) the trace of $A$ (resp. $B$), we have $\mathrm{Tr}(B) = \mathrm{Tr}(A) + 2c$. Since $\mathrm{Tr}(B) = -1$, we have

$$\mathrm{Tr}(A) \equiv -1 \bmod 2. \tag{4.4}$$

If $A$ is conjugate to the standard monodromy matrix of $I_2$, then $\mathrm{Tr}(A)$ is equal to 2, which contradicts (4.4). Thus the assertion is confirmed. □

- In the deformation (5.3) (see p.19), the type of the subordinate fiber is either (i) $II$, (ii) $I_2$ or (iii) $I_1 + I_1$. Now, by Proposition 4.6, we can eliminate the case (ii). Hence:

  $\underline{(5.3)\ IV \longrightarrow I_2 \text{ with } (II \text{ or } I_1 + I_1)}$

**Splittability of $II^*$**

$$II^* \to IV^* + II \qquad (s_0 s_2)^4 (s_0 s_2) = A_{IV^*} \cdot A_{II}$$

$$II^* \to I_2^* + I_1 + I_1 \qquad (s_0 s_2)^3 (s_0)^2 (s_0^{-1} s_2 s_0) s_2 = A_{I_2^*} \cdot A_{I_1}^{s_2} \cdot A_{I_1}^{s_0 s_2}$$

$$II^* \to I_5 + I_1 + I_1 + I_1 + I_1 + I_1$$

$$s_0^5 (s_0^{-1} s_2 s_0) s_0 s_2 s_2 s_0 = A_{I_5} \cdot A_{I_1}^{s_2} \cdot A_{I_1} \cdot A_{I_1}^{s_0 s_2} \cdot A_{I_1}^{s_0 s_2} \cdot A_{I_1}$$

$$II^* \to I_8 + I_1 + I_1 \qquad s_0^8 (s_0^{-2} s_2 s_0^2)(s_0^{-1} s_2^{-2} s_0 s_2^2 s_0) = A_{I_8} \cdot A_{I_1}^{s_0^{-1} s_2} \cdot A_{I_1}^{s_0^{-1} s_2^{-2}}$$

**Proposition 4.7.** *$II^*$ does not admit the following decompositions:*

$$II^* \longrightarrow I_8 + II$$

$$II^* \longrightarrow I_8 + I_2$$

*Proof.* This is showed by the same argument as that used in the proof of Proposition 4.6. □

- In the deformation (2.7) (see p.19), the type of the subordinate fiber is either (i) $II$, (ii) $I_2$ or (iii) $I_1 + I_1$. Now, by Proposition 4.7, we can eliminate the case (i) and (ii). Hence:

  $\underline{(2.7)\ II^* \longrightarrow I_8 \text{ with } I_1 + I_1}$

**Splittability of $III^*$**

$$\begin{array}{lcll}
III^* & \to & I_1^* + I_2 & s_2^{-1}(s_0s_2)^3 s_0 s_2 s_0^2 = A_{IV^*}^{s_2^{-1}} \cdot A_{I_2} \\
III^* & \to & I_0^* + I_1 + I_1 + I_1 & (s_0s_2)^3 s_0 s_2 s_0 = A_{I_0^*} \cdot A_{I_1} \cdot A_{I_1}^{s_0 s_2} \cdot A_{I_1} \\
III^* & \to & I_7 + I_1 + I_1 & s_0^7(s_0^{-5} s_2 s_0^5)(s_0^{-2} s_2 s_0^2) = A_{I_7} \cdot A_{I_1}^{s_0^{-4} s_2} \cdot A_{I_1}^{s_0^{-1} s_2} \\
III^* & \to & I_6 + II + I_1 & s_0^6(s_0^{-5} s_0 s_2 s_0^5)(s_0^{-2} s_2 s_0^2) = A_{I_6} \cdot A_{II}^{s_0^{-4} s_2} \cdot A_{I_1}^{s_0^{-1} s_2} \\
III^* & \to & I_6 + I_2 + I_1 & s_0^6(s_0^{-3} s_2 s_0^3)(s_2 s_0^2 s_2^{-1}) = A_{I_6} \cdot A_{I_1}^{s_0^{-2} s_2} \cdot A_{I_2}^{s_2} \\
III^* & \to & I_6 + I_1 + I_1 + I_1 &
\end{array}$$

**Proposition 4.8.** *$III^*$ does not admit the following decompositions:*

$$III^* \longrightarrow I_7 + II$$
$$III^* \longrightarrow I_7 + I_2$$
$$III^* \longrightarrow I_6 + III$$
$$III^* \longrightarrow I_6 + I_3$$

*Proof.* This is showed by the same argument as that used in the proof of Proposition 4.6. □

- In the deformation (4.7) (see p.19), the type of the subordinate fiber is either (i) $II$, (ii) $I_2$ or (iii) $I_1 + I_1$. Now, by Proposition 4.8, we can eliminate the case (i) and (ii). Hence:

  $\underline{(4.7)\ III^* \longrightarrow I_7 \text{ with } I_1 + I_1}$

- In the deformation (4.8) (see p.19), the type of the subordinate fiber is either (i) $III$, (ii) $I_3$, (iii) $II + I_1$, (iv) $I_2 + I_1$ or (v) $I_1 + I_1 + I_1$. Now, by Proposition 4.8, we can eliminate the case (i) and (ii). Hence:

  $\underline{(4.8)\ III^* \longrightarrow I_6 \text{ with } (II + I_1,\ I_2 + I_1 \text{ or } I_1 + I_1 + I_1)}$

**Splittability of $IV^*$**

$$\begin{array}{lcll}
IV^* & \to & I_0^* + I_1 + I_1 & (s_0s_2)^3 s_0 s_2 = A_{I_0}^* \cdot A_{I_1} \cdot A_{I_1}^{s_0 s_2} \\
IV^* & \to & I_6 + I_1 + I_1 & s_0^6(s_0^{-4} s_0 s_2 s_0^4)(s_0^{-1} s_2 s_0) = A_{I_6} \cdot A_{II}^{s_0^{-3} s_2} \cdot A_{I_1}^{s_2}
\end{array}$$

**Proposition 4.9.** *$IV^*$ does not admit the following decompositions:*

$$IV^* \longrightarrow I_6 + II$$
$$IV^* \longrightarrow I_6 + I_2$$

*Proof.* This is showed by the same argument as that used in the proof of Proposition 4.6. □

- In the deformation (6.3) (see p.19), the type of the subordinate fiber is either (i) $II$, (ii) $I_2$ or (iii) $I_1 + I_1$. Now, by Proposition 4.9, we can eliminate the case (i) and (ii). Hence:

  $\underline{(6.3)\ IV^* \longrightarrow I_6 \text{ with } I_1 + I_1}$

**Splittability of $I_0^*$**

$$\begin{array}{lll} I_0^* & \to & I_4 + I_1 + I_1 \qquad\qquad & s_0^4(s_0^{-2} s_2 s_0^2) s_2 = A_{I_4} \cdot A_{I_1}^{s_0^{-1} s_2} \cdot A_{I_1}^{s_0 s_2} \\ I_0^* & \to & I_3 + II + I_1 & s_0^3(s_0^{-2} s_0 s_2 s_0^2) s_2 = A_{I_3} \cdot A_{II}^{s_0^{-2} s_2} \cdot A_{I_1}^{s_0 s_2} \\ I_0^* & \to & I_3 + I_1 + I_1 + I_1 & \end{array}$$

**Proposition 4.10.** *$I_0^*$ does not admit the following decompositions:*

$$\begin{aligned} I_0^* &\longrightarrow I_4 + II \\ I_0^* &\longrightarrow I_4 + I_2 \\ I_0^* &\longrightarrow I_3 + III \\ I_0^* &\longrightarrow I_3 + I_3 \end{aligned}$$

*Proof.* If $I_0^*$ splits into $X_1 + X_2$, then for monodromy matrices $A_i$ $(i = 1, 2)$ of $X_i$, we have $-I = A_1 A_2$, that is,

$$A_1 = -A_2^{-1}.$$

In particular, $\mathrm{Tr}(A_1) = -\mathrm{Tr}(A_2)$. Noting that $\mathrm{Tr}(A_i)$ is determined by the type of $X_i$. this confirms the assertion. □

**Proposition 4.11.** *$I_0^*$ does not admit the following decomposition:*

$$I_0^* \longrightarrow I_3 + I_2 + I_1$$

*Proof.* If $I_0^*$ splits into $I_3 + X_1 + X_2$, then for monodromy matrices $A_i$ $(i = 1, 2)$ of $X_i$, we have $-I = I_3 A_1 A_2$. So

$$\begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix} A_1 = -A_2^{-1}.$$

In particular, writing $A_1 = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, $\mathrm{Tr}(A_1) + 3c = -\mathrm{Tr}(A_2)$ holds. Note that the traces of the standard monodromy matrices of $I_2$ and $I_1$ do not satisfy this condition. Thus the assertion is confirmed. □

- In the deformation (7.1) (see p.19), the type of the subordinate fiber is either (i) $II$, (ii) $I_2$ or (iii) $I_1 + I_1$. Now, by Proposition 4.10, we can eliminate the case (i) and (ii). Hence:

  $$\underline{(7.1)\ I_0^* \longrightarrow I_4 \text{ with } I_1 + I_1}$$

- In the deformation (7.2) (see p.19), the type of the subordinate fiber is either (i) $III$, (ii) $I_3$, (iii) $II + I_1$, (iv) $I_2 + I_1$ or (v) $I_1 + I_1 + I_1$. Now, by Propositions 4.10 and 4.11, we can eliminate the case (i) (ii) and (iv). Hence:

  $$\underline{(7.2)\ I_0^* \longrightarrow I_3 \text{ with } (II + I_1 \text{ or } I_1 + I_1 + I_1)}$$

**Splittability of $I_n^*$**

$$I_n^* \quad \to \quad I_{n+4} + I_1 + I_1 \qquad\qquad s_2 s_0^2 s_2 s_0^{-2} s_0^{n+4} = A_{I_1}^{s_0 s_2} \cdot A_{I_1}^{s_0^{-1} s_2} \cdot A_{I_{n+4}}$$

**Proposition 4.12.** *$I_n^*$ does not admit the following decompositions:*

$$I_n^* \longrightarrow I_{n+4} + II$$
$$I_n^* \longrightarrow I_{n+4} + I_2$$

*Proof.* This is showed by the same argument as that used in the proof of Proposition 4.6. □

- In the deformation (8.2) (see p.19), the type of the subordinate fiber is either (i) $II$, (ii) $I_2$ or (iii) $I_1 + I_1$. Now, by Proposition 4.12, we can eliminate the case (i) and (ii). Hence:

  (8.2) $I_n^* \longrightarrow I_{n+4}$ with $I_1 + I_1$

## 4.3 Singularities of subordinate fibers

We next consider the numbers of the subordinate fibers appearing in a barking family and their singularities. In fact, the upper bounds of them are determined by the data of the simple crust. Moreover if it satisfies some condition, then we may determine the numbers. Note that this observation depends on barking deformations, while that in the previous subsection does not.

Let $\pi : M \to \Delta$ be a linear degeneration of complex curves with a stellar singular fiber $X_0 = m_0\Theta_0 + \sum_{j=1}^h \mathrm{Br}^{(j)}$. If there exist a simple crust $Y$ of $X_0$, then we may construct a splitting family of $\pi : M \to \Delta$, which is called a barking family associated with $Y$ (see Theorem 2.6). Suppose that $Y = n_0\Theta_0 + \sum_{j=1}^h \mathrm{br}^{(j)}$ is a simple crust of $X_0$ with barking multiplicity $l$.

Recall that each subbranch of $Y$ is of type $A_l$, $B_l$ or $C_l$. A subbranch $\overline{\mathrm{br}}^{(j)}$ is said to be *proportional* if $m_0 n_1^{(j)} = n_0 m_1^{(j)}$ (equivalently $n_0/m_0 = n_1^{(j)}/m_1^{(j)} = \cdots = n_\nu^{(j)}/m_\nu^{(j)}$). Note that any proportional subbranch in simple crusts is of type $A_l$. Indeed, any proportional subbranch of type $B_l$ is of type $A_l$, and no proportional subbranch is of type $C_l$. Moreover any proportional subbranch $\overline{\mathrm{br}}^{(j)}$ has the same length as that of $\overline{\mathrm{Br}}^{(j)}$ (that is, $\nu^{(j)} = \lambda^{(j)}$).

The following result is important in order to investigate singularities of subordinate fibers ([Ta3] Proposition 16.2.6):

**Lemma 4.13.** *Suppose that $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$ is a barking family of the degeneration $\pi : M \to \Delta$ associated with a simple crust $Y$. Then any subordinate fiber of $\Psi$ is a reduced curve only with A-singularities*[5]*. Moreover these singularities lie* (a) *near the core or* (b) *near the edge of each proportional subbranch*[6] *if it exists.*

---

[5] An $A_\mu$-singularity is a singularity analytically equivalent to $y^2 = x^{\mu+1}$.

[6] Precisely speaking, the irreducible component (which is a projective line) at the end of the branch corresponding to each proportional subbranch.

**Remark 4.14.** Lemma 4.13 states that any subordinate fiber in barking families is a reduced curve only with isolated singularities, which ensures that none of $mI_n$ $(m \geq 2)$, $IV^*$, $III^*$, $II^*$ and $mI_n^*$ $(m \geq 2)$ appears as a subordinate fiber.

**Singularities near proportional subbranches** We will describe the singularities of subordinate fibers near a proportional subbranch. Let $\pi : M \to \Delta$ be a linear degeneration of complex curves with a stellar singular fiber $X_0$ and $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$ be a barking family associated with a simple crust $Y$ with barking multiplicity $l$. Suppose that $Y$ has a proportional subbranch br of a branch Br of $X_0$. First recall that near the branch Br, $\mathcal{M}$ is given by the following data (see [Ta3] Chapter 7): for $i = 1, 2, \dots, \lambda$,

$$\begin{cases} \mathcal{H}_i : w_i^{m_{i-1}-ln_{i-1}} \eta_i^{m_i - ln_i} (w_i^{n_{i-1}} \eta_i^{n_i} + t^d f_i)^l - s = 0, & \text{in } U_i \times \mathbb{C} \times \Delta \times \Delta^\dagger, \\ \mathcal{H}'_i : z_i^{m_{i+1}-ln_{i+1}} \zeta_i^{m_i - ln_i} (z_i^{n_{i+1}} \zeta_i^{n_i} + t^d \hat{f}_i)^l - s = 0, & \text{in } V_i \times \mathbb{C} \times \Delta \times \Delta^\dagger. \end{cases} \tag{4.5}$$

Note that substituting $t = 0$ into (4.5), then we obtain (2.2).

For fixed $(s, t) \in \Delta \times \Delta^\dagger$, we consider a fiber $X_{s,t} = \Psi^{-1}(s, t)$ of $\Psi$. The following is required:

**Lemma 4.15.** *Let $m, n, l$ be positive integers satisfying $m - ln > 0$ and $m', n'$ be nonnegative integers satisfying $m' - ln' \geq 0$. Set $h(z, \zeta) := f(z^{p'} \zeta^p)$ for a non-vanishing holomorphic function $f$ and positive integers $p, p'$ $(p < p')$. Then a complex curve $C_{s,t}$ in $\mathbb{C}^2$ defined by*

$$C_{s,t} \ : \ z^{m'-ln'} \zeta^{m-ln} (z^{n'} \zeta^n + th)^l - s = 0$$

*is singular if and only if*

(1) $s = 0$ *or*

(2) $m' = n' = 0$ *and* $\left(\dfrac{ln - m}{ln}\right)^{l\bar{n}} s^{\bar{n}} = \left(\dfrac{ln - m}{m} tc\right)^{\bar{m}}$,

*where $c := h(0, 0)$ and $\bar{m}$ and $\bar{n}$ are the relatively prime integers satisfying $\bar{n}/\bar{m} = n/m$. In the case (2), $(z, \zeta) \in C_{s,t}$ is a singularity exactly when $z = 0$ and $\zeta^n = \frac{ln-m}{m} tc$.*

Lemma 4.15 ensures that the hypersurface

$$\mathcal{H}'_\lambda\big|_{s,t} : \zeta_\lambda^{m_\lambda - ln_\lambda} (\zeta_\lambda^{n_\lambda} + t^d \hat{f}_\lambda)^l - s = 0, \quad \text{in } V_\lambda \times \mathbb{C} \times \{s\} \times \{t\},$$

has singularities, while the other hypersurfaces do not. In what follows, we write $m := m_\lambda$ and $n := n_\lambda$ for clarity.

For fixed $t \neq 0$, the equation $\left(\frac{ln-m}{ln}\right)^{l\bar{n}} s^{\bar{n}} = \left(\frac{ln-m}{m} t^d c\right)^{\bar{m}}$ for $s$ has $\bar{n}$ solutions, say $s_1, s_2, \dots, s_{\bar{n}}$. Noting that $(0, \zeta)$ satisfying $\zeta^n = \frac{ln-m}{m} tc$ is a singularity of $\mathcal{H}'_\lambda\big|_{s_k,t}$ for some $s_k$, each $\mathcal{H}'_\lambda\big|_{s_k,t}$ has $n/\bar{n}$ $(= \gcd(m, n))$ singularities respectively.

This result is summarized as follows:

**Proposition 4.16.** *Let $\pi : M \to \Delta$ be a linear degeneration of complex curves with a stellar singular fiber $X_0$ and let $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$ be a barking family associated with a simple crust $Y$ with barking multiplicity $l$. Suppose that $Y$ has a proportional subbranch $\mathrm{br}^{(j)}$ of a branch $\mathrm{Br}^{(j)}$ of $X_0$. Write $\overline{\mathrm{Br}}^{(j)} := m_0\Delta_0 + m_1\Theta_1 + m_2\Theta_2 + \cdots + m_\lambda\Theta_\lambda$ and $\overline{\mathrm{br}}^{(j)} := n_0\Delta_0 + n_1\Theta_1 + n_2\Theta_2 + \cdots + n_\lambda\Theta_\lambda$ and let $\bar{m}$ and $\bar{n}$ be the relatively prime integers satisfying $\bar{n}/\bar{m} = n_\lambda/m_\lambda$. Then for fixed $t \neq 0$, there exist $\bar{n}$ subordinate fibers that have singularities near the edge of $\mathrm{Br}^{(j)}$. Moreover, each subordinate fiber has $n/\bar{n}\,(= \gcd(m, n))$ singularities near the edge of $\mathrm{Br}^{(j)}$.*

**Singularities near the core** Next we will describe the singularities of subordinate fibers near the core.

Let $\pi : M \to \Delta$ be a linear degeneration of complex curves with a stellar singular fiber $X_0 = m_0\Theta_0 + \sum_{j=1}^h \mathrm{Br}^{(j)}$ and let $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$ be a barking family of the degeneration $\pi : M \to \Delta$ assosiated with a simple crust $Y = n_0\Theta_0 + \sum_{j=1}^h \mathrm{br}^{(j)}$. Write $\overline{\mathrm{Br}}^{(j)} = m_0\Delta_0^{(j)} + m_1^{(j)}\Theta_1^{(j)} + \cdots + m_{\lambda^{(j)}}^{(j)}\Theta_{\lambda^{(j)}}^{(j)}$, $\overline{\mathrm{br}}^{(j)} = n_0\Delta_0^{(j)} + n_1^{(j)}\Theta_1^{(j)} + \cdots + n_{\nu^{(j)}}^{(j)}\Theta_{\nu^{(j)}}^{(j)}$ and let $p^{(j)}$ be the attachment point on $\Theta_0$ with $\mathrm{Br}^{(j)}$. Recall that the local expression of $\mathcal{M}$ near the core $\Theta_0$ is given by

$$\sigma(z)\zeta^{m_0 - ln_0}\left(\zeta^{n_0} + t^d\tau(z)\right)^l - s = 0 \quad \text{in } N_0 \times \Delta \times \Delta^\dagger, \tag{4.6}$$

where $\sigma$ is the standard section of $N_0^{\otimes(-m_0)}$ on $\Theta_0$ and $\tau$ is a core section on $\Theta_0$ of $Y$. Substituting $t = 0$ into (4.6), then we obtain (2.1) for $i = 0$. Note that $\sigma$ has a zero of order $m_1^{(j)}$ at $p^{(j)}$, while $\tau$ has a pole of order $n_1^{(j)}$ at $p^{(j)}$. Suppose that $\tau$ has a zero of order $a_i$ at $q_i$ $(i = 1, 2, \ldots, k)$ on $\Theta_0$.

Fixing $s, t \neq 0$, consider a fiber $X_{s,t} := \Psi^{-1}(s, t)$ in $\Psi : \mathcal{M} \to \Delta \times \Delta^\dagger$. Set $F := \sigma(z)\zeta^{m_0 - ln_0}\left(\zeta^{n_0} + t^d\tau(z)\right)^l$. Then $(z, \zeta) \in X_{s,t}$ is a singularity precisely when $\frac{\partial}{\partial z}F(z, \zeta) = \frac{\partial}{\partial \zeta}F(z, \zeta) = 0$, equivalently

$$\begin{cases} \zeta^{m_0 - ln_0}(\zeta^{n_0} + t^d\tau(z))^{l-1}(\sigma_z(z)\zeta^{n_0} + t^d(\sigma_z(z)\tau(z) + l\sigma(z)\tau_z(z)) = 0, \\ \zeta^{m_0 - ln_0 - 1}(\zeta^{n_0} + t^d\tau(z))^{l-1}\sigma(z)(m_0\zeta^{n_0} + (m_0 - ln_0)t^d\tau_z(z)) = 0, \end{cases}$$

where $\sigma_z := \frac{d}{dz}\sigma$ and $\tau_z = \frac{d}{dz}\tau$. Setting $K(z) := n_0\sigma_z(z)\tau(z) + m_0\sigma(z)\tau_z(z)$[7], then we have

$$\begin{cases} K(z) = 0,\ \sigma(z) \neq 0,\ \tau(z) \neq 0 & (4.7) \\ \zeta^{n_0} = \dfrac{ln_0 - m_0}{m_0}t^d\tau(z). & (4.8) \end{cases}$$

Since points $(z, \zeta)$ in $X_{s,t}$ satisfy (4.6), substituting (4.8) into (4.6), we have

$$s = \left(\frac{ln_0}{ln_0 - m_0}\right)^l\sigma(z)\zeta^{m_0}.$$

Hence:

---

[7] Note that $K(z)$ is *not* a function on $\Theta_0$ but a meromorphic section of a line bundle $N_0^{\otimes(n-m)} \otimes \Omega^1_{\Theta_0}$ on $\Theta_0$, where $\Omega^1_{\Theta_0}$ is the cotangent bundle of $\Theta_0$.

**Lemma 4.17.** *Fix $t \neq 0$. A point $(z,\zeta) \in N_0$ is a singularity of some subordinate fiber $X_{s,t}$ if and only if the following is satisfied:*

$$\begin{cases} K(z) = 0,\ \sigma(z) \neq 0,\ \tau(z) \neq 0, & (4.9) \\ \zeta^{n_0} = \dfrac{ln_0 - m_0}{m_0} t^d \tau(z). & (4.10) \end{cases}$$

*In this case, the subordinate fiber $X_{s,t}$ satisfies*

$$s = (\frac{ln_0}{ln_0 - m_0})^l \sigma(z) \zeta^{m_0}. \tag{4.11}$$

We call a zero of $K(z)$ satisfying the condition (4.9) an *essential zero.* Combining (4.10) and (4.11), we have

$$s^{\bar{n}_0} = \left(\frac{ln_0}{ln_0 - m_0}\right)^{l\bar{n}_0} \sigma(z)^{\bar{n}_0} \left(\frac{ln_0 - m_0}{m_0}\right)^{\bar{m}_0} t^{d\bar{m}_0} \tau(z)^{\bar{m}_0}, \tag{4.12}$$

where $\bar{m}_0$ and $\bar{n}_0$ are the relatively prime integers satisfying $\bar{n}_0/\bar{m}_0 = n_0/m_0$. For an essential zero $\alpha$ of $K(z)$, clearly this equation for $s$ has $\bar{n}_0$ solutions, say $s_1, s_2, \ldots, s_{\bar{n}_0}$. On the other hand, the equation (4.10) for $\zeta$ has $n_0$ solutions, say $\beta_1, \beta_2, \ldots, \beta_{n_0}$, which implies that $n_0/\bar{n}_0 = \gcd(m_0, n_0)$ points among $(\alpha, \beta_1), (\alpha, \beta_2), \ldots, (\alpha, \beta_{n_0})$ lies on one of subordinate fibers $X_{s_1,t}, X_{s_2,t}, \ldots, X_{s_{\bar{n}_0},t}$.

**Lemma 4.18.** *Let $\alpha$ be an essential zero of $K(z)$. Then there exist $\bar{n}_0$ subordinate fibers that have a singularity with $z$-coordinate $\alpha$. Moreover the number of such singularities of each of them is $n_0/\bar{n}_0$.*

Next we express $K(z) = \sigma\tau\omega$, where $\omega(z) := \dfrac{d\log(\sigma^{n_0}\tau^{m_0})}{dz}$. Here $\omega$ is a meromorphic section of the cotangent bundle $\Omega^1_{\Theta_0}$ on $\Theta_0$. For $\omega(z)$, the following holds:

(1) $\omega(z)$ has a pole of order 1 at $p^{(j)}$ if $m_0 n_1^{(j)} - m_1^{(j)} n_0 \neq 0$, whereas $\omega(z)$ is holomorphic at $p^{(j)}$ if $m_0 n_1^{(j)} - m_1^{(j)} n_0 = 0$.

(2) $\omega(z)$ has a pole of order 1 at $q_i$ for $i = 1, 2, \ldots, k$.

Set $J_0 := \left\{ j \mid m_0 n_1^{(j)} - m_1^{(j)} n_0 = 0 \right\} \subset \{1, 2, \ldots, h\}$ (so $\overline{\mathrm{br}}^{(j)}$ is a proportional subbranch for $j \in J_0$). Then

$$\begin{cases} K(z) = 0, \\ \sigma(z) \neq 0, \\ \tau(z) \neq 0 \end{cases} \iff \begin{cases} \omega(z) = 0, \\ z \neq p_j\ (j \in J_0). \end{cases}$$

**Lemma 4.19.** *Set $v := \#J_0$ and denote by $g_0$ the genus of $\Theta_0$. Then*

$$\sum_{K(\alpha)=0, \sigma(\alpha)\neq 0, \tau(\alpha)\neq 0} \mathrm{ord}_\alpha(K(z)) = (h - v) + k + (2g_0 - 2) - \sum_{j \in J_0} \mathrm{ord}_{p^{(j)}}(\omega).$$

We set $\chi := (h-v)+k+(2g_0-2)-\sum_{j\in J_0}\mathrm{ord}_{p^{(j)}}(\omega)$, which is called the *core invariant*. From Lemma 4.19, we have:

**Corollary 4.20.**

$$\big(\textit{the number of essential zeros of } K(z)\big) \le \chi,$$

*where the equality holds precisely when the order of any essential zero equals* 1.

Where $\kappa$ is the number of essential zeros of $K(z)$, let $\alpha_1, \alpha_2, \dots, \alpha_\kappa$ be the essential zeros of $K(z)$. We denote by $\bar{\kappa}$ the number of the distinct values of the set $\{\sigma(\alpha_j)^{\bar{n}_0}\tau(\alpha_j)^{\bar{m}_0} \mid j = 1,2,\dots,\kappa\}$. Note that if $\alpha_i$ and $\alpha_j$ satisfy

$$\sigma(\alpha_i)^{\bar{n}_0}\tau(\alpha_i)^{\bar{m}_0} = \sigma(\alpha_j)^{\bar{n}_0}\tau(\alpha_j)^{\bar{m}_0},$$

then the equations (4.12) for $s$ in the cases $z=\alpha_i$ and $z=\alpha_j$ coincide, so that they have the same $\bar{n}_0$ solutions. Thus, for fixed $t\neq 0$, there exist $\bar{n}_0\bar{\kappa}$ subordinate fibers that have singularities near the core. This result together with Lemma 4.18 and Corollary 4.20 confirms the following:

**Proposition 4.21.** *For fixed $t \neq 0$,*

1. $\big(\textit{the number of subordinate fibers with singularities near } \Theta_0\big) \le \bar{n}_0\chi$, *where the equality holds precisely when the order of any essential zero equals* 1 *and* $\bar{\kappa}=\kappa$.
2. $\big(\textit{the number of singularities near } \Theta_0 \textit{ on each subordinate fiber}\big) \le \dfrac{n_0}{\bar{n}_0}\chi$, *where the equality holds precisely when the order of any essential zero equals* 1 *and* $\bar{\kappa}=1$.

**The case that $Y$ has no proportional subbranch** Now we return to the grobal observation. Let $\pi : M \to \Delta$ be a linear degeneration of complex curves with a stellar singular fiber $X_0 = m_0\Theta_0 + \sum_{j=1}^h \mathrm{Br}^{(j)}$. Suppose that $X_0$ has a simple crust $Y = n_0\Theta_0 + \sum_{j=1}^h \mathrm{br}^{(j)}$ of with barking multiplicity $l$. Let $\Psi : \mathcal{M} \to \Delta\times\Delta^\dagger$ be a barking family of $\pi : M\to\Delta$ associated with $Y$. We define the core invariant $\chi$ of $Y$ by

$$\chi := (h-v)+k+(2g_0-2)-\sum_{j\in J_0}\mathrm{ord}_{p^{(j)}}(\omega),$$

where $J_0 := \left\{ j \;\middle|\; m_0 n_1^{(j)} - m_1^{(j)} n_0 = 0 \right\} \subset \{1,2,\dots,h\}$, $v := \#J_0$ (that is, the number of proportional subbranches of $Y$), $\omega := \frac{d}{dz}\log(\sigma^{n_0}\tau^{m_0})$ and $g_0$ is the genus of $\Theta_0$ (see Lemma 4.19).

First we consider the case that $Y$ has no proportional subbranch. Then Lemma 4.13 ensures that singularities of subordinate fibers lies only near the core. Note that $\chi = h+k+(2g_0-2)$.

**Lemma 4.22.** *Suppose that $Y$ has no proportional subbranch. Set $c := \gcd(m_0, n_0)$ and $\bar{n}_0 := n_0/c$. If $\chi = 1$, then for fixed $t\neq 0$, the following holds:*

(1) *$\pi_t : M_t \to \Delta$ contains $\bar{n}_0$ subordinate fibers.*

(2) *Each subordinate fiber in $\pi_t : M_t \to \Delta$ has $c$ singularities.*

(3) *The number of singularities of all the subordinate fibers in $\pi_t : M_t \to \Delta$ is $n_0$.*

*Proof.* From Corollary 4.20, $\big(\text{the number of essential zeros of } K(z)\big) \leq \chi$. Since $K(z)$ has at least a zero, by assumption, we have

$$\big(\text{the number of essential zeros of } K(z)\big) = 1,$$

which implies that $K(z)$ has just one zero of order 1. Then from Proposition 4.21, we have

$$\begin{aligned}&\big(\text{the number of subordinate fibers in } \pi_t : M_t \to \Delta\big) = \bar{n}_0,\\ &\big(\text{the number of singularities on each subordinate fiber in } \pi_t : M_t \to \Delta\big) = c,\end{aligned}$$

confirming (1) and (2). Clearly (3) follows from (1) and (2). □

In particular:

**Proposition 4.23.** *Suppose that* (a) *$\Theta_0$ is a projective line,* (b) *$X_0$ has three branches,* (c) *the core section $\tau$ has no zero and* (d) *$Y$ has no proportional subbranch. Set $c := \gcd(m_0, n_0)$ and $\bar{n}_0 := n_0/c$. Then for fixed $t \neq 0$, the following holds:*

(1) *$\pi_t : M_t \to \Delta$ contains $\bar{n}_0$ subordinate fibers.*

(2) *Each subordinate fiber in $\pi_t : M_t \to \Delta$ has $c$ singularities.*

(3) *The number of singularities of all the subordinate fibers in $\pi_t : M_t \to \Delta$ is $n_0$.*

*Proof.* By assumption, we have $g_0 = 0$, $h = 3$, $k = 0$, and so $\chi = 1$. It is immediate from Lemma 4.22. □

**Remark 4.24.** By Lemma 2.5, the condition (c) is restated as "$r_0 = r'_0$ holds," where $r_0 := \sum_{j=1}^{h} m_1^{(j)}/m_0$ and $r'_0 := \sum_{j=1}^{h} n_1^{(j)}/n_0$.

- In the deformations (3.2) and (4.2) (see p.19), the type of the subordinate fiber is either (i) $II$, (ii) $I_2$ or (iii) $I_1 + I_1$. Now, applying Proposition 4.23, it is easy to see that there appears just one subordinate fiber which has two singularities. Hence:

  $\underline{(3.2)\ III \longrightarrow I_1 \text{ with } I_2}$

  $\underline{(4.2)\ III^* \longrightarrow I_1^* \text{ with } I_2}$

- In the deformations (5.2) and (6.2) (see p.19), the type of the subordinate fiber is either (i) $II$, (ii) $I_2$ or (iii) $I_1 + I_1$. Now, applying Proposition 4.23, it is easy to see that there appear two subordinate fibers each of which has one singularity. Hence:

$$\underline{(5.2)\ IV \longrightarrow I_2 \text{ with } I_1 + I_1}$$
$$\underline{(6.2)\ IV^* \longrightarrow I_0^* \text{ with } I_1 + I_1}$$

- In the deformations (4.4) and (4.5) (see p.19), the type of the subordinate fiber is either (i) $III$, (ii) $I_3$, (iii) $II + I_1$, (iv) $I_2 + I_1$ or (v) $I_1 + I_1 + I_1$. Now, applying Proposition 4.23, it is easy to see that there appear three subordinate fibers each of which has one singularity. Hence:

$$\underline{(4.4)\ III^* \longrightarrow I_0^* \text{ with } I_1 + I_1 + I_1}$$
$$\underline{(4.5)\ III^* \longrightarrow I_6 \text{ with } I_1 + I_1 + I_1}$$

- In the deformation (2.4) (see p.20), applying Proposition 4.23, it is easy to see that there appear five subordinate fibers each of which has one singularity. Hence:

$$\underline{(2.4)\ II^* \longrightarrow I_5 \text{ with } I_1 + I_1 + I_1 + I_1 + I_1}$$

**The case that $Y$ has a proportional subbranch** Next we assume that $Y$ has a proportional subbranch.

**Proposition 4.25.** *Suppose that* (a) $\Theta_0$ *is a projective line,* (b) $X_0$ *has three branches,* (c) *the core section* $\tau$ *has no zero and* (d) $Y$ *has a proportional subbranch* $\overline{\mathrm{br}}^{(1)} = n_0\Delta_0 + n_1\Theta_1 + n_2\Theta_2 + \cdots + n_\lambda\Theta_\lambda$ *of* $\overline{\mathrm{Br}}^{(1)}$. *Then* $\overline{\mathrm{br}}^{(1)}$ *is the unique proportional subbranch of* $Y$. *Moreover for fixed* $t \neq 0$, *the following holds:*

(1) $\pi_t : M_t \to \Delta$ *contains* $\bar{n}_\lambda$ *subordinate fibers.*

(2) *Each subordinate fiber in* $\pi_t : M_t \to \Delta$ *has* $c$ *singularities.*

(3) *The number of singularities of all the subordinate fibers in* $\pi_t : M_t \to \Delta$ *is* $n_\lambda$.

*Here* $c := \gcd(m_\lambda, n_\lambda)$ *and* $\bar{n}_\lambda := n_\lambda / c$.

*Proof.* By assumption, we have $g_0 = 0$, $h = 3$, $k = 0$. Thus

$$\begin{aligned} \chi &= 1 - v - \sum_{j \in J_0} \mathrm{ord}_{p^{(j)}}(\omega) \\ &= 1 - \sum_{j \in J_0} \left( \mathrm{ord}_{p^{(j)}}(\omega) + 1 \right). \end{aligned}$$

Recall that $\omega(z)$ is holomorphic at $p^{(j)}$ for $j \in J_0$ (that is, $m_0 n_1^{(j)} - m_1^{(j)} n_0 = 0$), which means that $\mathrm{ord}_{p^{(j)}}(\omega) \geq 0$. Since $\chi \geq 0$, we deduce $\mathrm{ord}_{p^{(j)}}(\omega) = 0$ and

$v = 1$. Hence $\overline{\mathrm{br}}^{(1)}$ is the unique proportional subbranch. Noting that $\chi = 0$, from Proposition 4.21, any subordinate fiber in $\pi_t : M_t \to \Delta$ has no singularity near the core $\Theta_0$ but near the edge of $\overline{\mathrm{Br}}^{(1)}$. Therefore Propositon 4.16 confirms the assertion. □

- In the deformation (2.2) (see p.19), the type of the subordinate fiber is either (i) $II$, (ii) $I_2$ or (iii) $I_1 + I_1$. Now, applying Proposition 4.25, it is easy to see that there appears just one subordinate fiber which has one singularity. Hence:

  $$\underline{(2.2)\ II^* \longrightarrow IV^* \text{ with } II}$$

- In the deformation (2.3) (see p.19), the type of the subordinate fiber is either (i) $II$, (ii) $I_2$ or (iii) $I_1 + I_1$. Now, applying Proposition 4.25, it is easy to see that there appears two subordinate fibers each of which has one singularity. Hence:

  $$\underline{(2.3)\ II^* \longrightarrow I_2^* \text{ with } I_1 + I_1}$$

# References

[BPV] W. Barth, C. Peter, and Van de Ven, Compact Complex Surfaces, Ergebnisse der Mathematik und ihrer Grentzgebiete (3. Folge), 4, Springer, 1984.

[FM] R. Friedman and J. Morgan, Smooth Four-Manifolds and Complex Surfaces, Ergebnisse der Mathematik und ihrer Grentzgebiete (3. Folge), 27, Springer, 1994.

[Ka] A. Kas, On the deformation types of regular elliptic surfaces, Complex Analysis and Algebraic Geometry, Iwanami shoten publ. and Cambridge Univ. Press, (1977), 107–112.

[Ko] K. Kodaira, On compact complex analytic surfaces II, Ann. of Math. 77, (1963), 563–626.

[M] B. Moishezon, Complex Surfaces and Connected Sums of Complex Projective Planes, Lecture Notes in Math. 603, Springer, 1977.

[Ta1] S. Takamura, Towards the classification of atoms of degenerations, I, (Splitting criteria via configurations of singular fibers), Journal of the Mathematical Society of Japan, 56 (1), (2004), 115–145.

[Ta2] S. Takamura, Towards the classification of atoms of degenerations, II - linearization of degenerations of complex curves, RIMS Preprint, 1344, (2001).

[Ta3] S. Takamura, Towards the Classification of Atoms of Degenerations, III - Splitting Deformations of Degenerations of Complex Curves, Lecture Notes in Math. 1886, Springer, 2006.

[U] M. Ue, On the Topology of Elliptic Surfaces, a Surgey, Amer. Math. Soc. Trans. (2) 160, (1994), 95–123.

GRADUATE SCHOOL OF MATHEMATICS, KYUSHU UNIVERSITY, MOTOOKA 744, NISHI-KU, FUKUOKA 819-0395, JAPAN
*E-mail address*: `t-okuda@math.kyushu-u.ac.jp`